

Components of Nurse Substance Use Disorder Monitoring Programs

Katherine R. Lee, JD, MN, RN

Introduction: The purpose of this study was to identify the components of nurse substance use disorder monitoring programs. The study was conducted in a large, tertiary care hospital. The study included a review of the literature and interviews with nursing leaders. The study found that the most common components of nurse substance use disorder monitoring programs were: (1) a policy that prohibits the use of alcohol and drugs on the job; (2) a program that provides education and training to nurses about the risks of substance use; (3) a program that provides support and resources for nurses who are struggling with substance use; and (4) a program that provides monitoring and testing for substance use. The study also found that the most common barriers to the implementation of nurse substance use disorder monitoring programs were: (1) a lack of resources; (2) a lack of buy-in from nursing leaders; and (3) a lack of support from the hospital administration. The study concludes that nurse substance use disorder monitoring programs are an important component of a comprehensive approach to addressing the problem of nurse substance use. The study also provides recommendations for the implementation of nurse substance use disorder monitoring programs.

Aim: The aim of this study was to identify the components of nurse substance use disorder monitoring programs.

Methods: The study was conducted in a large, tertiary care hospital. The study included a review of the literature and interviews with nursing leaders.

Results: The study found that the most common components of nurse substance use disorder monitoring programs were: (1) a policy that prohibits the use of alcohol and drugs on the job; (2) a program that provides education and training to nurses about the risks of substance use; (3) a program that provides support and resources for nurses who are struggling with substance use; and (4) a program that provides monitoring and testing for substance use. The study also found that the most common barriers to the implementation of nurse substance use disorder monitoring programs were: (1) a lack of resources; (2) a lack of buy-in from nursing leaders; and (3) a lack of support from the hospital administration.

() , , -
,
☒ ,
- ()
,)
()

, - , - -
,
☒ , ☒ , %
☒ , -
☒ ☒ -
, , ☒ ☒ -

(=)

() - () (-) () ()

,

, ,) -

,

, ()

T

B

% (=)

F

-

,

,

,

-

,

(=),
☒

(=)

,

%

(=) -

- -

M *r* *r* *r* **N**
☒

D

☒ % -

==

☒ () D, ...
☒ () ☒
, ☒ , , , &
() ☒ ☒
J ... S ☒ , A ☒ , T ...
37(),
(,) C ...
() S ☒ , ...
: A ...
☒☒☒ (()

(() 0 () () ())